

## LA-UR-20-23791

Approved for public release; distribution is unlimited.

Title: Impact of a minority relativistic electron tail interacting with a thermal plasma containing high-atomic-number impurities

Author(s): Garland, Nathan Ashley

Intended for: Online presentation to ITER collaboration. Will be shared after presentation on their internal share site

Issued: 2020-05-21

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



# Impact of a minority relativistic electron tail interacting with a thermal plasma containing high-atomic-number impurities

Nathan Garland (LANL), Hyun-Kyung Chung (NFRI, Korea), Chris Fontes<sup>A</sup>, Mark Zammit<sup>A</sup>, James Colgan<sup>A</sup>, Todd Elder, Chris McDevitt, Tim Wildey (SNL), Xianzhu Tang

ITER DMS Task Force - 4<sup>th</sup> May 2020

UNCLASSIFIED

# Impact of a minority relativistic electron tail interacting with a thermal plasma containing high-atomic-number impurities

Cite as: Phys. Plasmas **27**, 040702 (2020); doi: [10.1063/5.0003638](https://doi.org/10.1063/5.0003638)

Submitted: 4 February 2020 · Accepted: 7 April 2020 ·

Published Online: 24 April 2020 · Corrected: 28 April 2020

[View Online](#)[Export Citation](#)[CrossMark](#)

Nathan A. Garland,<sup>1,a)</sup> Hyun-Kyung Chung,<sup>2</sup> Christopher J. Fontes,<sup>1</sup> Mark C. Zammit,<sup>1</sup> James Colgan,<sup>1</sup> Todd Elder,<sup>1,3</sup> Christopher J. McDevitt,<sup>1,4</sup> Timothy M. Wildey,<sup>5</sup> and Xian-Zhu Tang<sup>1</sup>

## AFFILIATIONS

<sup>1</sup>Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

<sup>2</sup>National Fusion Research Institute (NFRI), 169-148 Gwahak-ro, Yuseong-gu, Daejeon 34133, South Korea

<sup>3</sup>Columbia University, New York, New York 10027, USA

<sup>4</sup>Department of Materials Science and Engineering, University of Florida, Gainesville, Florida 32611, USA


<sup>5</sup>Sandia National Laboratories, Albuquerque, New Mexico 87185, USA

<sup>a)</sup>Author to whom correspondence should be addressed: [ngarland@lanl.gov](mailto:ngarland@lanl.gov)


UNCLASSIFIED

# Motivation: Tokamak disruptions

- Current ITER strategy: inject high Z impurities to terminate (Ne, Ar)
  - Dissipate magnetic energy via radiation
  - Regulate plasma current
  - *But how does a plasma with runaway electrons interact with these impurities?*



RE impacting carbon tiles on Tore Supra  
(Source: CEA-IRFM)



Cooled molten Be tiles in JET  
(Source: EUROfusion)


Ionization balance/charge state population?

Radiation?


UNCLASSIFIED

# Motivation: Tokamak disruptions

- Current ITER strategy: inject high Z impurities to terminate (Ne,Ar)
  - Dissipate magnetic energy via radiation
  - Regulate plasma current
  - *But how does a plasma with runaway electrons interact with these impurities?*



RE impacting carbon tiles on Tore Supra  
(Source: CEA-IRFM)



Cooled molten Be tiles in JET  
(Source: EUROfusion)

Ionization balance/charge state population?

Radiation?



Collisional-Radiative modeling

UNCLASSIFIED

# CR models in the fusion community


- US groups have long used KPRAD to couple with plasma codes and probe experiments on DIII-D, C-MOD [Whyte *et al.* Proceedings of the 24th European Conference on Controlled Fusion and Plasma Physics 21A:1137 (1997)]
- EU groups have used ADAS at a mature level [Summers *et al.* AIP Conference Proceedings 901, no. 1 (2007)]
- HEDP often uses FLYCHK [Chung *et al.* HEDP 1 3 (2005)]

## General idea of this work:

*Do we need to do anything different from standard modeling to describe impact of runaways, and does it make a difference?*

UNCLASSIFIED


# Interaction of EEDF with inelastic cross-sections



UNCLASSIFIED



# Interaction of EEDF with inelastic cross-sections



UNCLASSIFIED

# Our approach

- Developing fork of FLYCHK [Chung *et al.* HEDP 13 (2005)], flychklite, to allow relativistic corrections & arbitrary EEDF
- QED effects manifest at relativistic  $e^-$  energies, generalized Breit interaction [Fontes *et al.* PRA **47** 2 (1993)]
  - Higher order correction to Coulomb interaction
  - Generally employed for binding energy, but not always cross-sections
- Enhanced  $e^-e^-$  coupling increases cross-section

UNCLASSIFIED

# Required input data: relativistic ICS

- We modify base inelastic ICS in FLYCHK to transition to a relativistic approximate ICS

$$\sigma_{i \rightarrow j}^{\text{TOT}} = (1 - S(E))\sigma_{i \rightarrow j}^{\text{NR}} + S(E)\sigma_{i \rightarrow j}^{\text{R}}$$

- Simulate relativistic effect via Møller-Bethe-like analytic relativistic-rise

$$\sigma_I^{\text{rel}} \sim \left( \log\left(\frac{\beta^2}{1 - \beta^2} \frac{0.5m_e c^2}{\Delta I_Z^i}\right) - \beta^2 \right)$$

$$\beta = v/c$$

$$\frac{dn_i}{dt} = -n_i \sum_{j \neq i}^{N_L} W_{ij} + \sum_{j \neq i}^{N_L} n_j W_{ji} \quad 1 \leq i \leq N_L$$

## Model assumptions:

In this work solving, SS problem

Optically thin


No IPD

UNCLASSIFIED

# Simulate relativistic effect via Møller-Bethe-like analytic form

- Møller-Bethe-like form not a new thing
- Back to 1930s
  - Moller *Annalen der Physik* 406 5 (1932)
  - Bethe *Z. Physik* 76 5 (1932)
- Provides a general prescription we can apply to any inelastic collision
  - But we try to benchmark against QM calculations

## Ar L-shell ionization ICS



FLYCHK/Burgess & Chidichimo MNRAS 203 1269 (1983)

Scofield PRA 18 963 (1978)

Wang *et al.* JPB 51 145201 (2018)



UNCLASSIFIED

# What effect does this have?

UNCLASSIFIED

# D+Ar plasma with 10 MeV Gaussian tail


$n_D = n_{Ar} = 10^{14} \text{ cm}^{-3}$ ; Runaway fraction  $\sim 10^{10}$ - $10^{11} \text{ cm}^{-3}$   
carrying an ITER-like current ( $\sim 10 \text{ MA}$ )



UNCLASSIFIED

# Ar CSD spread with cooling $T_e$

$$T_e = 10000.0 eV$$





- CSD spread as plasma cools is crucial to  $e^-$  scattering
- Runaway tail yields greater mean charge, over a wide range of charged states

UNCLASSIFIED

# D+Ar plasma with ITER-like currents

$n_D = n_{Ar} = 10^{14} \text{ cm}^{-3}$ ; Runaway fraction set to carry current  $\sim 10^{10} - 10^{11} \text{ cm}^{-3}$ .




ATOMIC - LANL CR model [Fontes *et al.* JPB 48 144014 (2015)]

UNCLASSIFIED



# Is this effect real? UQ can help



Ranges computed from epistemic UQ study with Dakota software



[\[https://dakota.sandia.gov/\]](https://dakota.sandia.gov/)

Vary relativistic rise of  $\sigma_{\text{ion}}$  and  $\sigma_{\text{exc}}$  with prefactors between 0.5 and 2, to account for variation in approximate formula

500 samples over parameter space with Latin Hypercube Sampling.

UNCLASSIFIED

# CSD effect on Bethe stopping power



$$\text{BSP: } -\frac{dE}{dx} = \sum_{\alpha} \frac{e^2}{\epsilon_0 m_e c^2} \frac{n_{e\alpha}}{\beta^2} \left[ \ln \left( \frac{2m_e c^2 \beta^2}{\langle I_{\alpha} \rangle (1 - \beta^2)} \right) - \beta^2 \right]$$


■ Thermal CSD  
● RA enhanced CSD

UNCLASSIFIED


# CSD effect on friction & deflection collision frequencies

Using formulae presented in Hesslow *et al.* PRL118 25 (2017)  
Compared with Rosenbluth & Putvinski, NF 37, 1355 (1997)


Normalized coll freq. 3.899 eV



Normalized coll freq. 12.824 eV



Normalized coll freq. 68.962 eV



$$\nu_S^{ee} = \nu_{S,cs}^{ee} \left[ 1 + \sum_j \frac{n_j N_{e,j}}{n_e \ln \Lambda} \left( \frac{1}{k} \ln(1 + h_j^k) - \beta^2 \right) \right] \quad \nu_D^{ei} = \nu_{D,cs}^{ei} \left( 1 + \frac{1}{Z_{\text{eff}}} \sum_j \frac{n_j g_j(p)}{n_e \ln \Lambda} \right)$$

UNCLASSIFIED

# Summary

- Including enhanced relativistic ICS → potential for enhanced  $\langle Z \rangle$  and RPL at lower  $T_e$  when minority RE are present
  - These low  $T_e$  are crucial for cooled, post-disruption discharges
- We must continue exploring
  - RE energies, profiles, densities
  - Current machines as well as ITER
  - Improving atomic physics input data

UNCLASSIFIED


# Where to from here?

- Time-dependent cooling history
  - Rapid timescales in disruptions, how bad are steady-state data approximations?
- Couple CR into plasma codes for consistent evolution of runaway EEDF,  $T_e$ , impurity ions
  - Continuum RFP model
  - 6D particle code (Chris McDevitt)
- Experimental verification
  - Discussing with tokamak operators
  - Controlled lab experiment

UNCLASSIFIED

# Where to from here?

- Further understand impact of RE and relativistic ICS to discharge composition
  - unique spectra produced by runaways
  - Joint effort of simpler flychklite CR with detailed LANL ATOMIC code
- Impact of excitations



(Left) Charge state with excitations, and (Right) without excitations (coronal assumption)

UNCLASSIFIED